TRINITY PRESBYTERIAN NURSERY SCHOOL FOOD ALLERGY EMERGENCY ACTION PLAN

	FOOD ALLERGY EMERGE	ENCY ACTION PLAN		
CHILD'S NA	ME:	D.O.B//	Child's	
CLASS:	TWOs TTH3s MWF3s _	MWF Pre-Ks PM Pre-Ks	Photograph	
SEVERE	ALLERGY TO:			
MILD AL	LERGY TO:			
ASTHMA	(higher risk for severe reaction)		Weight:lbs	
	f checked, give epinephrine immediately f checked, give epinephrine immediately			
SEVERE SYMPTOMS		INJECT EPINEP	INJECT EPINEPHRINE IMMEDIATELY	
ONE OR MORE OF THE FOLLOWING: LUNGS Short of breath, wheezing, repetitive cough HEART Pale, blue, faint, weak pulse, dizzy THROAT Tight, hoarse, trouble breathing/swallowing MOUTH Obstructive swelling (tongue and/or lips) SKIN Many hives over body OR A COMBINATION OF SYMPTOMS: SKIN Hives, itchy rashes, swelling (e.g., eyes, lips) GUT Vomiting, crampy pain		Pring Pring BEGIN MONIT Note time epin Give inhaler / I When in doubt, use	BEGIN MONITORING (see below) Note time epinephrine was administered	
MILD SYM	PTOMS ONLY	GIVE ANTIHIST	AMINE	
ONE OR MOUTH - SKIN - GUT	ORE OF THE FOLLOWING: Itchy mouth A few hives around mouth / face, mild it Mild nausea/discomfort	Stay with child professionals a	; alert healthcare and parent S PROGRESS, INJECT	
ambulance wit or more after t	child; alert healthcare professionals and the child; alert healthcare professionals and the child; he child; he first if symptoms persist or recur. For a subtild even if parents cannot be reached.	was administered. A second dose ma	y be administered 5 minutes	
MEDICATION	S			
EPINEPHRINI		Dose		
ANTHIHISTAN				
OTHER (e.g. in	haler / bronchilator) Brand			
EMERGENCY	CONTACTS			
Parent/Guardian:		Phone:		
Parent/Guardian:				
Name/Relationship:				
Name/Relationship:				

Physician/Healthcare Provider

Date

Parent/Guardian Signature

Date